

## 兔抗 SLC22A6 多克隆抗体

中文名称: 兔抗 SLC22A6 多克隆抗体

英文名称: Anti-SLC22A6 rabbit polyclonal antibody

别 名: OAT1, PAHT, HOAT1, ROAT1

相关类别: 一抗

储 存: 冷冻(-20℃)

宿 主: Rabbit

抗 原: SLC22A6

反应种属: Human, Rat

标 记 物: Unconjugate

克隆类型: rabbit polyclonal

## 技术规格

**Background:** 

The protein encoded by this gene is involved in the sodium-dependent transport and excret ion of organic anions, some of which are pot entially toxic. The encoded protein is an integ ral membrane protein and may be localized t o the basolateral membrane. Four transcript v ariants encoding four different isoforms have been found for this gene. Functions as organi c anion exchanger when the uptake of one m olecule of organic anion is coupled with an ef flux of one molecule of endogenous dicarbox ylic acid (glutarate, ketoglutarate, etc). Mediat es the sodium-independent uptake of 2,3-dim



## www.shjning.com

|                             | ercapto-1-propanesulfonic acid (DMPS) By sim ilarity.          |
|-----------------------------|--|
| Applications:               | ELISA, WB, IHC   |
| Name of antibody:           | SLC22A6  |
| Immunogen:                  | Synthetic peptide of human SLC22A6                             |
| Full name:                  | Solute carrier family 22 (organic anion transporter), member 6 |
| Synonyms:                   | OAT1, PAHT, HOAT1, ROAT1                                       |
| SwissProt:                  | Q4U2R8   |
| ELISA Recommended dilution: | 2000-10000   |
| IHC positive control:       | Human colon cancer and Human liver cancer                      |
| IHC Recommend dilution:     | 50-200   |
| WB Predicted band size:     | 56 kDa   |
| WB Positive control:        | Human fetal kidney tissue                                      |
| WB Recommended dilution:    | 500-2000   |





