

兔抗 MED28 多克隆抗体

中文名称: 兔抗 MED28 多克隆抗体

英文名称: Anti-MED28 rabbit polyclonal antibody

别 名: mediator complex subunit 28; EG1; magicin; 1500003D12Rik

相关类别: 一抗

储 存: 冷冻 (-20℃)

宿 主: Rabbit

抗 原: MED28

反应种属: Human, Mouse, Rat

标 记 物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

Applications: ELISA, IHC	participates in cellular signaling to the actin cytoskeleton downstream of tyrosine kinase signaling pathways.	rs. May be part of a complex containing NF2/merlin that	Background:	ved in the regulated transcription of nearly all RNA poly merase II-dependent genes. Mediator functions as a brid ge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription ma chinery. Mediator is recruited to promoters by direct inte ractions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex wi th RNA polymerase II and the general transcription facto rs. May be part of a complex containing NF2/merlin that participates in cellular signaling to the actin cytoskeleton downstream of tyrosine kinase signaling pathways.
participates in cellular signaling to the actin cytoskeleton	,		Background:	th RNA polymerase II and the general transcription facto
rs. May be part of a complex containing NF2/merlin that participates in cellular signaling to the actin cytoskeleton	rs. May be part of a complex containing NF2/merlin that	th RNA polymerase II and the general transcription facto		for the assembly of a functional preinitiation complex wi
th RNA polymerase II and the general transcription facto rs. May be part of a complex containing NF2/merlin that participates in cellular signaling to the actin cytoskeleton	th RNA polymerase II and the general transcription factors. May be part of a complex containing NF2/merlin that	·		ractions with regulatory proteins and serves as a scaffold
ractions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex wi th RNA polymerase II and the general transcription facto rs. May be part of a complex containing NF2/merlin that participates in cellular signaling to the actin cytoskeleton	ractions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex wi th RNA polymerase II and the general transcription facto rs. May be part of a complex containing NF2/merlin that	ractions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex wi		chinery. Mediator is recruited to promoters by direct inte
ractions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex wi th RNA polymerase II and the general transcription factors. May be part of a complex containing NF2/merlin that participates in cellular signaling to the actin cytoskeleton	ractions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors. May be part of a complex containing NF2/merlin that	ractions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex wi		proteins to the basal RNA polymerase II transcription ma
chinery. Mediator is recruited to promoters by direct inte ractions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex wi th RNA polymerase II and the general transcription factors. May be part of a complex containing NF2/merlin that participates in cellular signaling to the actin cytoskeleton	chinery. Mediator is recruited to promoters by direct inte ractions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors. May be part of a complex containing NF2/merlin that	Chinery. Mediator is recruited to promoters by direct inte ractions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex wi		ge to convey information from gene-specific regulatory
Background: Background: proteins to the basal RNA polymerase II transcription ma chinery. Mediator is recruited to promoters by direct inte ractions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex wi th RNA polymerase II and the general transcription factors. May be part of a complex containing NF2/merlin that participates in cellular signaling to the actin cytoskeleton	Background: Background: proteins to the basal RNA polymerase II transcription ma chinery. Mediator is recruited to promoters by direct inte ractions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex wi th RNA polymerase II and the general transcription factors. May be part of a complex containing NF2/merlin that	proteins to the basal RNA polymerase II transcription ma chinery. Mediator is recruited to promoters by direct inte ractions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex wi		merase II-dependent genes. Mediator functions as a brid
ge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription ma chinery. Mediator is recruited to promoters by direct inte ractions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex wi th RNA polymerase II and the general transcription facto rs. May be part of a complex containing NF2/merlin that participates in cellular signaling to the actin cytoskeleton	ge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription ma chinery. Mediator is recruited to promoters by direct inte ractions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex wi th RNA polymerase II and the general transcription factors. May be part of a complex containing NF2/merlin that	ge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription ma chinery. Mediator is recruited to promoters by direct inte ractions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex wi		ved in the regulated transcription of nearly all RNA poly
merase II-dependent genes. Mediator functions as a brid ge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription ma chinery. Mediator is recruited to promoters by direct inte ractions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex wi th RNA polymerase II and the general transcription factors. May be part of a complex containing NF2/merlin that participates in cellular signaling to the actin cytoskeleton	merase II-dependent genes. Mediator functions as a brid ge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription ma chinery. Mediator is recruited to promoters by direct inte ractions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex wi th RNA polymerase II and the general transcription factors. May be part of a complex containing NF2/merlin that	merase II-dependent genes. Mediator functions as a brid ge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription ma chinery. Mediator is recruited to promoters by direct inte ractions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex wi		Component of the Mediator complex, a coactivator invol



Name of antibody:	MED28
Immunogen:	Fusion protein of human MED28
Full name:	mediator complex subunit 28
Synonyms:	EG1; magicin; 1500003D12Rik
SwissProt:	Q9H204
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human liver cancer
IHC Recommend dilution:	200-400

