

## 兔抗 PLAUR 多克隆抗体

- 中文名称：兔抗 PLAUR 多克隆抗体
- 英文名称：Anti-PLAUR rabbit polyclonal antibody
- 别名：CD87; UPAR; URKR; U-PAR
- 抗原：PLAUR
- 储存：冷冻（-20℃）
- 宿主：Rabbit
- 相关类别：一抗
- 反应种属：Human
- 标记物：Unconjugate
- 克隆类型：rabbit polyclonal

### 技术规格

**Background:**

This gene encodes the receptor for urokinase plasminogen activator and, given its role in localizing and promoting plasmin formation, likely influences many normal and pathological processes related to cell-surface plasminogen activation and localized degradation of the extracellular matrix. It binds both the proprotein and mature forms of urokinase plasminogen activator and permits the activation of the receptor-bound pro-enzyme by plasmin. The protein lacks transmembrane or cytoplasmic domains and may be anchored to the plasma membrane by a glycosyl-phosphatidylinositol (GPI) moiety following cleavage of the nascent polypeptide near its carboxy-terminus. However, a soluble protein is also produced in some cell types. Altern

	ative splicing results in multiple transcript variants encoding different isoforms. The proprotein experiences several post-translational cleavage reactions that have not yet been fully defined.
<b>Applications:</b>	ELISA, WB, IHC
<b>Name of antibody:</b>	PLAUR
<b>Immunogen:</b>	Fusion protein of human PLAUR
<b>Full name:</b>	Plasminogen activator, urokinase receptor
<b>Synonyms :</b>	CD87; UPAR; URKR; U-PAR
<b>SwissProt:</b>	Q03405
<b>ELISA Recommended dilution:</b>	2000-5000
<b>IHC positive control:</b>	Human lung cancer and Human colon cancer
<b>IHC Recommend dilution:</b>	50-200
<b>WB Predicted band size:</b>	37 kDa
<b>WB Positive control:</b>	A549 cells
<b>WB Recommended dilution:</b>	500-2000



